arXiv:1212.4766v1 [math-ph] 19 Dec 2012

Contractions of 2D 2nd order quantum superintegrable systems and the Askey scheme for hypergeometric orthogonal polynomials

Ernest G. Kalnins,
Department of Mathematics, University of Waikato,
Hamilton, New Zealand,
math0236@waikato.ac.nz
Willard Miller, Jr.
School of Mathematics, University of Minnesota,
Minneapolis, Minnesota, 55455, U.S.A.
miller@ima.umn.edu
and Sarah Post
Department of Mathematics, U. Hawaii at Manoa
Honolulu HI 96822
spost@hawaii.edu

Abstract

We show explicitly that all 2nd order superintegrable systems in 2 dimensions are limiting cases of a single system: the generic 3-parameter potential on the 2-sphere, $S9$ in our listing. Analogously we show that all of the quadratic symmetry algebras of these systems are contractions of that of $S9$. By contracting function space realizations of irreducible representations of the S9 algebra (which give the structure equations for Racah/Wilson polynomials) to the other superintegrable systems we obtain the full Askey scheme of orthogonal hypergeometric polynomials. This relationship directly ties the polynomials and their structure equations to physical phenomena. It is more general because it applies to all special functions that arise from these systems via separation of variables, not just those of hypergeometric type, and it extends to higher dimensions.

1 Introduction

A quantum superintegrable system is an integrable Hamiltonian system on an n-dimensional Riemannian/pseudo-Riemannian manifold with potential: $H = \Delta_n + V$ that admits $2n-1$ algebraically independent partial differential operators commuting with H, apparantly the maximum possible. Superintegrability captures the properties of quantum Hamiltonian systems that allow the Schrödinger eigenvalue problem $H\Psi = E\Psi$ to be solved exactly, analytically and algebraically, see e.g., [1, 2, 3, 4, 5, 6, 7, 8, 9, 10]. A system is of order N if the maximum order of the symmetry operators, other than H, is N. For $n = 2$, $N = 2$ all systems are known. There are about 50 types on a variety of manifolds, [11] II, but under the Stäckel transform, an invertible structure preserving mapping, they divide into 12 equivalence classes with representatives on flat space and the 2-sphere, 6 with nondegenerate 3-parameter potentials and 6 with degenerate 1-parameter potentials, [12], [13]. The symmetry operators of each system close to generate a quadratic algebra, and the irreducible representations of this algebra determine the eigenvalues of H and their multiplicity [14, 15, 16, 17, 18, 19, 20]. All the 2nd order superintegrable systems are limiting cases of a single system: the generic 3-parameter potential on the 2-sphere, $S9$ in our listing. Analogously all quadratic symmetry algebras of these systems are contractions of $S9$. There are also contractions of $S9$ to systems that admit 3 independent symmetry operators and relate to the Askey Scheme but are such that the metric becomes singular. We refer to these systems as "singular".

The irreducible representations of $S9$ have a realization in terms of difference operators in 1 variable, exactly the structure algebra for the Wilson and Racah polynomials! By contracting these representations to obtain the representations of the quadratic symmetry algebras of the other superintegrable systems we obtain the full Askey scheme of orthogonal hypergeometric polynomials, [21, 22]. This relationship ties the structure equations directly to physical phenomena. For example, for the Meixner-Pollaczek and Pseudo Jacobi polynomials and special cases of the Wilson, continuous Hahn and continuous dual Hahn polynomials, there are instances where the parameters in these functions occur in complex conjugate pairs but a real three term recurrence relation exists, so that the polynomials are orthogonal with respect to a positive weight function. In these instances, the quantum Hamiltonian is *PT-symmetric* in the sense that the scalar potential $V(x, y)$ satisfies $V(x, y) = \overline{V(y, x)}$. Here T means time inversion (complex conjugation) and P means permutation. When the Hamiltonian admits PT symmetry then even though the potential is complex, the bound state energy eigenvalues must be real. PT symmetry in physics is controversial but the mathematics is clear [23]. The usual meaning of P in physics is space inversion, so that PT symmetry requires $V(x,y) = \overline{V(-x,-y)}$, but the outcome is the same: the potential is complex but the energy eigenvalues are real. The superintegrable system $S6$, an analog of the 2D hydrogen atom on the 2-sphere, is an example of PT symmetry in the standard sense, [2] (2012).

The method relates structure equations; pointwise polynomial limits are just a byproduct, it applies to all special functions that arise from these systems via separation of variables, not just polynomials of hypergeometric type, and it extends to higher dimensions, [24] (2011). In some cases, e.g. for Hahn polynomials, we have contractions leading to structure equations not obeyed by the full family of Hahn polynomials. These families with higher than usual symmetry we refer to as "special". The structure theory exposes these "special" systems in a natural manner.

There is a close association with the symmetry operators in the original physical quantum systems that describe bases of eigenfunctions via separation of variables. All of the quantum systems are multiseparable. Some separable systems are exactly solvable in the sense that the physical solutions are products of hypergeometric functions. The special functions arising in the models we consider can be described as the coefficients in the expansion of a separable eigenbasis for the original quantum system in terms of another separable eigenbasis. The functions in the Askey Scheme are all hypergeometric polynomials that arise as the expansion coefficients relating two separable eigenbases that are *both* of hypergeometric type. Nondegenerate systems $E2$, $E10$ and degenerate system $E4$, defined below, fail this test, so they are not involved with the Askey scheme, though they fit the contraction scheme and describe other special functions. In $E2$ separation occurs in two distinct eigenbases, but only one of is exactly solvable; for $E10$ separation occurs in two eigenbases, but neither is exactly solvable; for $E4$ there is separation in two exactly solvable systems but the expansion coefficients are not hypergeometric.

Now we provide more detail about 2D 2nd order superintegrable systems. In local coordinates x_i the Hamiltonian takes the form $H = \Delta_2 + V(\mathbf{x})$ where $\Delta_2 = \frac{1}{\sqrt{g}} \sum_{ij=1}^{2} \partial_i (g^{ij} \sqrt{g} \partial_j)$ is the Laplace-Beltrami operator in these coordinates, $g^{ij}(\mathbf{x})$ is the contravariant metric tensor and g is the determinant of the covariant metric tensor. A 2nd order symmetry operator for this system is a partial differential operator $L = \frac{1}{\sqrt{g}} \sum_{ij=1}^{2} \partial_i (S^{ij}(\mathbf{x}) \sqrt{g} \partial_j) + W(\mathbf{x})$, where S^{ij} is a symmetric contravariant tensor, such that $[H,S] \equiv HS - SH = 0$. (These operators are formally self-adjoint with respect to the bilinear product $< f, g >_\lambda = \int f(\mathbf{x}) g(\mathbf{x}) \lambda(\mathbf{x}) dx_1 dx_2$ on the manifold,[24] (2010). The system is *2nd order superintegrable* if there are two symmetry operators S_1, S_2

such that the set $\{H, L_1, L_2\}$ is algebraically independent, i.e., there is no nontrivial polynomial $P(H, L_1, L_2)$, symmetric in L_1, L_2 such that $P \equiv 0$. Note that if there is only one symmetry operator L_1 then the system is 2nd order integrable. The requirement that two symmetry operators exist is highly restrictive. It turns out for our treatment of 2nd order 2D quantum superintegrable systems the values of the mass m and Planck's constant $\hbar$ are immaterial, so we have normalized our Hamiltonians as given.

Since every 2D Riemannian space is conformally flat, we can always assume the existence of Cartesian-like coordinates x_1, x_2 such that

$$H = \frac{1}{\lambda(\mathbf{x})}(\partial_{11} + \partial_{22}) + V(\mathbf{x}),\ L_k = \frac{1}{\lambda}\sum_{i,j=1}^{2} \partial_i(L^{ij}_{(k)}\lambda\partial_j) + W_{(k)}(x\mathbf{x}),\ k = 1, 2.$$

The commutators $[H, L_k] = 0$, $k = 1, 2$ put conditions on the the potentials $W_{(1)}, W_{(2)}$, enabling us to solve for the partial derivatives $\partial_i W_{(k)}$ in terms of the function V and its 1st derivatives. The integrability conditions $\partial_1(\partial_2 W_{(k)}) = \partial_2(\partial_1 W_{(k)})$, the Bertrand-Darboux equations [11] I, lead to the necessary and sufficient condition that V must satisfy a pair of coupled linear equations of the form

$$V_{22} - V_{11} = A^{22}V_1 + B^{22}V_2, \quad V_{12} = A^{12}V_1 + B^{12}V_2, \tag{1}$$

for locally analytic functions $A^{ij}(\mathbf{x}), B^{ij}(\mathbf{x})$. Here $V_i = \partial_i V$, etc. We call these the canonical equations. If the integrability equations for (1) are satisfied identically then the solution space for the canonical equations is 4-dimensional and we can always express the general solution in the form $V(\mathbf{x}) = a_1V_{(1)}(\mathbf{x}) + a_2V_{(2)}(\mathbf{x}) + a_3V_{(3)}(\mathbf{x}) + a_4$ where a_4 is a trivial additive constant. In this case we say that the potential is *nondegenerate* and refer to it as 3-parameter. Another possibility is that the solution space is 2-dimensional with general solution $V(\mathbf{x}) = a_1V_{(1)}(\mathbf{x}) + a_2$. In this case we say that the potential is *degenerate* and refer to it as 1-parameter. Every degenerate potential can be obtained from some nondegenerate potential by restricting the parameters. It is not *just* a restriction, however, because the structure of the symmetry algebra changes. A formally skew-adjoint 1st order symmetry appears and this induces a new 2nd order symmetry. The last possibility is that the integrability conditions are satisfied only by a constant potential. In that case we refer to the system as *free*; the equation $H\Psi = E\Psi$ is just the Laplace-Beltrami eigenvalue equation. The case of a two-parameter potential doesn't occur, i.e., any 2-parameter potential extends to a 3-parameter potential, [13].

All of these systems have the remarkable property that the symmetry algebras generated by H, S_1, S_2 for nondegenerate potentials close under commutation. Define the 3rd order commutation R by $R = [S_1, S_2]$. Then the fourth order operators $[R, S_1], [R, S_2]$ are contained in the associative algebra of symmetrized products of the generators [11] I:

$$[L_j, R] = \sum_{0\le e_1+e_2+e_3\le 2} M^{(j)}_{e_1,e_2,e_3}\{L_1^{e_1}, L_2^{e_2}\}H^{e_3}, \qquad e_k \ge 0,$$

where $\{L_1, L_2\} = L_1L_2 + L_2L_1$ is the symmetrizer. Also the 6th order operator R^2 is contained in the algebra of symmetrized products up to 3rd order:

$$R^2 - \sum_{0\le e_1+e_2+e_3\le 3} N_{e_1,e_2,e_3}\{L_1^{e_1}, L_2^{e_2}\}H^{e_3} = 0. \tag{2}$$

In both equations the constants $M^{(j)}_{e_1,e_2,e_3}$ and N_{e_1,e_2,e_3} polynomials in the parameters a_1, a_2, a_3 of degree $2-e_1-e_2-e_3$ and $3-e_1-e_2-e_3$ respectively.

For systems with one parameter potentials the situation is different [13]. There are 4 generators: one 1st order X and 3 second order H, L_1, L_2. The commutators $[X, S_1], [X, S_2]$ are 2nd order and expressed as

$$[X, S_j] = \sum_{0\le e_1+e_2+e_3+e_4\le 1} P^{(j)}_{e_1,e_2,e_3,e_4} L_1^{e_1}L_2^{e_2}H^{e_3}X^{e_4}, \quad j = 1, 2. \tag{3}$$

The commutator $[L_1, L_2]$ is 3rd order skew adjoint and expressed as

$$[L_1, L_2] = \sum_{0\le e_1+e_2+e_3+e_4\le 1} Q_{e_1,e_2,e_3,e_4}\{L_1^{e_1}L_2^{e_2}, X\}H^{e_3}X^{2e_4}.$$

Finally, since there are at most 3 algebraically independent generators, there must be a polynomial identity satisfied by the 4 generators. It is of 4th order:

$$G \equiv \sum_{0\le e_1+e_2+e_3+e_4\le 2} S_{e_1,e_2,e_3,e_4}\{L_1^{e_1}, L_2^{e_2}, X^{2e_4}\}H^{e_3}, \tag{4}$$

where $\{L_1^{e_1}, L_2^{e_2}, X^{2e_4}\}$ is the symmetrizer of three operators and has 6 terms. The constants $P^{(j)}_{e_1,e_2,e_3,e_4}$, Q_{e_1,e_2,e_3,e_4} and S_{e_1,e_2,e_3,e_4} are polynomials in the parameter a_1 of degrees $1-e_1-e_2-e_3-e_4$, $1-e_1-e_2-e_3-e_4$ and $2-e_1-e_2-e_3-e_4$ respectively.

All of the possibilities have been classified. The classification is simplified greatly by use of the Stäckel transform, an invertible structure preserving mapping from a superintegrable system on one manifold to a superintegrable system on another manifold,[24] (2010). Thus, if we know the structure equations for H then we know the structure equations for any system Stäckel equivalent to H. For our study we can restrict ourselves to a choice of one representative system in each equivalence class. There are 13 Stäckel equivalence classes of systems with nonfree potentials but one is an isolated Euclidean singleton unrelated to the Askey scheme. In [11] II, it is shown that every 2nd order 2D superintegrable system is Stäckel equivalent to a constant curvature system, so we will choose our examples in flat space and on complex 2-spheres. In [25] all 21 such systems in flat space are determined up to equivalence under the complex Euclidean group and all 9 nonzero constant curvature spaces are determined up to equivalence under the complex orthogonal group. (Some of these systems are Stäckel equivalent to one another.) We will use the notation given there. There are 6 Stäckel equivalence classes of nondegenerate potentials and 6 of degenerate potentials.

2 The 12 superintegrable systems

In this section, we fix some notation. Let $s_1^2 + s_2^2 + s_3^2 = 1$ be the embedding of the 2-sphere in 3D Euclidean space and $z = x + iy$, $\overline{z} = x - iy$. Define $J_3 = s_1\partial_{s_2} - s_2\partial_{s_1}$ to be the generator of rotation about the s_3 axis, with J_2, J_3 obtained by cyclic permutation. On the Euclidean plane, we shall also use J_3 to denote the generator of rotations in the x, y plane. As in the previous section $R = [L_1, L_2]$.

1) Quantum S9: This quantum superintegrable system is defined,

$$\begin{aligned} H =& \quad J_1^2 + J_2^2 + J_3^2 + \tfrac{a_1}{s_1^2} + \tfrac{a_2}{s_2^2} + \tfrac{a_3}{s_3^2}, \\ L_1 =& \quad J_1^2 + \tfrac{a_3 s_2^2}{s_3^2} + \tfrac{a_2 s_3^2}{s_2^2} \\ L_2 =& \quad J_2^2 + \tfrac{a_1 s_3^2}{s_1^2} + \tfrac{(a_3 s_1^2}{s_3^2}. \end{aligned} \tag{5}$$

The algebra is given by

$$\begin{aligned} [L_i, R] =& \, 4\{L_i, L_k\} - 4\{L_i, L_j\} - (8 + 16a_j)L_j + (8 + 16a_k)L_k + 8(a_j - a_k), \\ R^2 =& \, \tfrac{8}{3}\{L_1, L_2, L_3\} - (16a_1 + 12)L_1^2 - (16a_2 + 12)L_2^2 - (16a_3 + 12)L_3^2 \\ & + \tfrac{52}{3}(\{L_1, L_2\} + \{L_2, L_3\} + \{L_3, L_1\}) + \tfrac{1}{3}(16 + 176a_1)L_1 \\ & + \tfrac{1}{3}(16 + 176a_2)L_2 + \tfrac{1}{3}(16 + 176a_3)L_3 + \tfrac{32}{3}(a_1 + a_2 + a_3) \\ & + 48(a_1a_2 + a_2a_3 + a_3a_1) + 64a_1a_2a_3. \end{aligned} \tag{6}$$

In practice we make the substitution $L_3 = H - L_1 - L_2 - a_1 - a_2 - a_3$, so that the generators are H, L_1, L_2.

2) Quantum E1: The quantum system is defined by

$$\begin{aligned} H =& \quad \partial_x^2 + \partial_y^2 + a_1(x^2 + y^2) + \tfrac{a_2}{x^2} + \tfrac{a_3}{y^2} \\ L_1 =& \quad \partial_y^2 + a_1 y^2 + \tfrac{a_3}{y^2} \\ L_2 =& \quad (x\partial_y - y\partial_x)^2 + \left(\tfrac{a_2 y^2}{x^2} + \tfrac{a_3 x^2}{y^2}\right). \end{aligned} \tag{7}$$

The algebra relations are

$$\begin{aligned}[L_1,R] =&\ 8L_1H-8L_1^2+16a_1L_2-8a_1(1+2a_2+2a_3)\\ [L_2,R] =&\ 8\{L_1,L_2\}-4L_2H-16(1+a_2+a_3)L_1+8(1+2a_3)H\\ R^2 =&\ \tfrac{8}{3}\left(\{L_1,L_2,H\}-\{L_1,L_1,L_2\}\right)-(16a_3+12)H^2\\ &-\left(\tfrac{176}{3}+16a_2+16a_3\right)L_1^2-16a_1L_2^2+\left(\tfrac{176}{3}+32a_3\right)L_1H\\ &+\tfrac{176a_1}{3}L_2-\tfrac{16a_1}{3}\left(12a_2a_3+9a_2+9a_3+2\right).\end{aligned} \tag{8}$$

3) Quantum E2: The generators are

$$\begin{aligned}H =&\ \partial_x^2+\partial_y^2+a_1(4x^2+y^2)+a_2x+\tfrac{a_3}{y^2},\\ L_1 =&\ \partial_y^2+a_1y^2+\tfrac{a_3}{y^2},\\ L_3 =&\ \tfrac{1}{2}\{(x\partial_y-y\partial_x),\partial_y\}-y^2(a_1x-\tfrac{a_2}{4})+\tfrac{a_3x}{y^2}.\end{aligned} \tag{9}$$

$$\begin{aligned}[L_1,R] =&\ 2a_2L_2+16a_1L_2,\\ [L_2,R] =&\ -2L_1^2+4L_2^2-4L_2H+2a_2L_3-a_1(8a_3+6),\\ R^2 =&\ 4L_2^2+4L_1H+16a_1L_2^2-2a_2\{L_1,L_2\}+(12+16a_3)a_1L_1\\ &+32a_1^2L_3-a_2^2(a_3+\tfrac{3}{4}).\end{aligned} \tag{10}$$

4) Quantum E3′: The quantum system is defined by

$$\begin{aligned}H =&\ \partial_x^2+\partial_y^2+a_1(x^2+y^2)+a_2x+a_3y+\tfrac{a_2^2+a_3^2}{4a_1},\\ L_1 =&\ \partial_y^2+a_1y^2+a_3y+\tfrac{a_3^2}{4a_1}.\\ L_2 =&\ 2\partial_x\partial_y+\tfrac{2(a_1x+a_2)(2a_1y+a_3)}{2a_1},\end{aligned} \tag{11}$$

with algebra relations

$$\begin{aligned}[L_1,R] =&\ -4a_1L_2,\\ [L_2,R] =&\ 16a_1L_1+8a_1H\\ R^2 =&\ 16a_1L_1H-16a_1L_1^2-4a_1L_2^2-16a_1^2.\end{aligned} \tag{12}$$

5) Quantum E10: The quantum system is defined by

$$\begin{aligned}H =&\ 4\partial\overline{\partial}+a_1\left(z\overline{z}-\tfrac{1}{2}\overline{z}^3\right)+a_2\left(z-\tfrac{3}{2}\overline{z}^2\right)+a_3\overline{z}\\ L_1 =&\ -\partial^2-\tfrac{a_1\overline{z}^2}{4}-\tfrac{a_2\overline{z}}{2}+\tfrac{a_3}{12}\\ L_2 =&\ \{z\partial-\overline{z}\overline{\partial},\partial\}-\overline{\partial}^2-(2z+\overline{z}^2)\left(\tfrac{a_1(2z-3\overline{z}^2)}{16}-\tfrac{a_2\overline{z}}{2}+\tfrac{a_3}{4}\right).\end{aligned}$$

The algebra relations are

$$\begin{aligned}[L_1,R] =&\ 2a_1L_1-\tfrac{a_2^2}{2}-\tfrac{a_1a_3}{6}\\ [L_2,R] =&\ 24L_1^2+4a_3L_1-2a_1L_2+a_2H\\ R^2 =&\ -16L_1^3-\tfrac{a_1}{4}H^2+2a_1\{L_1,L_2\}-2a_2L_1H-4a_3L_1^2\\ &-(a_2^2+\tfrac{a_1a_3}{3})L_2-\tfrac{a_2a_3}{3}H-a_1^2+\tfrac{a_3^3}{27}.\end{aligned} \tag{13}$$

6) Quantum E8: The quantum system is defined by $(\partial = \partial_z)$:

$$\begin{aligned} H = & \ 4\partial\overline{\partial} + a_1 z\overline{z} + \tfrac{a_2 z}{\overline{z}^3} + \tfrac{a_3}{\overline{z}^2} \\ L_1 = & \ -\partial^2 - \tfrac{a_1}{4}\overline{z}^2 + \tfrac{a_2}{2\overline{z}^2}, \\ L_2 = & \ -\left(z\partial + \overline{z}\overline{\partial}\right)^2 + \tfrac{a_2 z^2}{\overline{z}^2} + \tfrac{a_3 z}{\overline{z}}. \end{aligned} \tag{14}$$

The algebra relations are:

$$\begin{aligned} & [L_1, R] = -8L_1^2 + 2a_1a_2 \\ & [L_2, R] = 8\{L_1, L_2\} - 16L_1 - 2a_3H \\ & R^2 = 8\{L_1^2, L_2\} - \tfrac{176}{3}L_1^2 - a_3L_1H + a_2H^2 - 4a_1a_2L_2 - \tfrac{a_1(3a_3^2 - 4a_2)}{3}. \end{aligned} \tag{15}$$

2.1 Degenerate quantum systems

7) Quantum S3 (Higgs oscillator): The system is the same as $S9$ with $a_1 = a_2 = 0$ $a_3 = a$. The symmetry algebra is generated by

$$X = J_3, \qquad L_1 = J_1^2 + \frac{as_2^2}{s_3^2}, \qquad L_2 = \frac{1}{2}(J_1J_2 + J_2J_1) - \frac{as_1s_2}{s_3^2}.$$

The structure relations for the algebra are given by

$$\begin{aligned} & [L_1, X] = 2L_2, \qquad [L_2, X] = -X^2 - 2L_1 + H - a, \\ & [L_1, L_2] = -(L_1X + XL_1) - (\tfrac{1}{2} + 2a)X, \\ & 0 = \{L_1, X^2\} + 2L_1^2 + 2L_2^2 - 2L_1H + \tfrac{5+4a}{2}X^2 - 2aL_1 - a. \end{aligned} \tag{16}$$

8) Quantum E14: The system is defined by

$$\begin{aligned} & H = \partial_x^2 + \partial_y^2 + \tfrac{a}{\overline{z}^2}, & X = \partial \\ & L_1 = \tfrac{i}{2}\{z\partial + \overline{z}\overline{\partial}, \partial\} + \tfrac{a}{\overline{z}}, & L_2 = (z\partial + \overline{z}\overline{z})^2 + \tfrac{az}{\overline{z}}, \end{aligned} \tag{17}$$

with structure equations

$$\begin{aligned} & [L_1, L_2] = -\{X, L_2\} - \tfrac{1}{2}X, & [X, L_1] = -X^2, \\ & [X, L_2] = 2L_1, & L_1^2 + XL_2X - bH - \tfrac{1}{4}X^2 = 0. \end{aligned} \tag{18}$$

9) Quantum E6: The system is defined by

$$\begin{aligned} & H = \partial_x^2 + \partial_y^2 + \tfrac{a}{x^2}, & X = \partial_y \\ & L_1 = \tfrac{1}{2}\{x\partial_y - y\partial_x, \partial_x\} - \tfrac{ay}{x^2}, & L_2 = (x\partial_y - y\partial_x)^2 + \tfrac{ay^2}{x^2}, \end{aligned} \tag{19}$$

with symmetry algebra

$$\begin{aligned} & [L_1, L_2] = \{X, L_2\} + (2a + \tfrac{1}{2})X, \quad [L_1, X] = H - X^2, \quad [L_2, X] = 2L_1, \\ & L_1^2 + \tfrac{1}{4}\{L_2, X^2\} + \tfrac{1}{2}XL_2X - L_2H + (a + \tfrac{3}{4})X^2 = 0. \end{aligned} \tag{20}$$

10) Quantum E5: The system is defined by

$$\begin{array}{ll} H = \partial_x^2 + \partial_y^2 + ax, & X = \partial_y \\ L_1 = \partial_{xy} + \frac{1}{2}ay, \qquad L_2 = \frac{1}{2}\{x\partial_y - y\partial_x, \partial_y\} - \frac{1}{4}ay^2. & \end{array} \tag{21}$$

The structure equations are

$$\begin{array}{l} [L_1, L_2] = 2X^3 - HX, \qquad [L_1, X] = -\frac{a}{2}, \qquad [L_2, X] = L_1, \\ X^4 - HX^2 + L_1^2 + aL_2 = 0. \end{array} \tag{22}$$

11) Quantum E4: The system is defined by

$$\begin{array}{ll} H = \partial_x^2 + \partial_y^2 + a(x + iy), & X = \partial_x + i\partial_y \\ L_1 = \partial_x^2 + ax, & L_2 = \frac{i}{2}\{x\partial_y - y\partial_x, X\} - \frac{a}{4}(x + iy)^2. \end{array} \tag{23}$$

The structure equations are

$$\begin{array}{c} [L_1, X] = a, \quad [L_2, X] = X^2, \quad [L_1, L_2] = X^3 + HX - \{L_1, X\}, \\ X^4 - 2\{L_1, X^2\} + 2HX^2 + H^2 + 4aL_2 = 0. \end{array} . \tag{24}$$

12) Quantum E3(harmonic oscillator): The system is determined by

$$\begin{array}{c} H = \partial_x^2 + \partial_y^2 + a(x^2 + y^2), \quad X = x\partial_y - y\partial_x \\ L_1 = \partial_y^2 - \omega^2 y^2, \quad L_2 = \partial_{xy} + axy. \end{array}$$

The structure equations are

$$\begin{array}{c} [L_1, X] = 2L_2, \quad [L_2, X] = H - 2L_1, \quad [L_1, L_2] = -2aX \\ L_1^2 + L_3^2 - L_1 H + aX^2 - a = 0. \end{array} \tag{25}$$

2.2 Free (1st order) Quantum superintegrable systems

1) The 2-sphere: Here $s_1^2 + s_2^2 + s_3^2 = 1$ is the embedding of the 2-sphere in Euclidean space, and the Hamiltonian is $H = J_1^2 + J_2^2 + J_3^2$, where $J_3 = s_1\partial_{s_2} - s_2\partial_{s_1}$ and J_2, J_3 are obtained by cyclic permutations of $1, 2, 3$. The basis symmetries are J_1, J_2, J_3. They generate the Lie algebra $so(3)$ with relations $[J_1, J_2] = -J_3$, $[J_2, J_3] = -J_1$, $[J_3, J_1] = -J_2$ and Casimir H.

2) The Euclidean plane: Here $H = \partial_x^2 + \partial_y^2$ with basis symmetries $P_1 = \partial_x$, $P_2 = \partial_y$ and $M = x\partial_y - y\partial_x$. The symmetry Lie algebra is $e(2)$ with relations $[P_1, P_2] = 0$, $[P_1, M] = P_2$, $[P_2, M] = -P_1$ and Casimir H.

3 Contractions of superintegrable systems

We will give a detailed treatment of contractions in another publication [26], but here we just describe "natural" contractions. Suppose we have a nondegenerate superintegrable system with generators H, L_1, L_2 and structure equations (2), defining a quadratic algebra Q. If we make a change of basis to new generators $\tilde{H}, \tilde{L}_1, \tilde{L}_2$ and parameters $\tilde{a}_1, \tilde{a}_2, \tilde{a}_3$ such that

$$\begin{pmatrix} \tilde{L}_1 \\ \tilde{L}_2 \\ \tilde{H} \end{pmatrix} = \begin{pmatrix} A_{1,1} & A_{1,2} & A_{1,3} \\ A_{2,1} & A_{2,2} & A_{2,3} \\ 0 & 0 & A_{3,3} \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \end{pmatrix} + \begin{pmatrix} B_{1,1} & B_{1,2} & B_{1,3} \\ B_{2,1} & B_{2,2} & B_{2,3} \\ B_{3,1} & B_{3,2} & B_{3,3} \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix},$$

$$\begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix} = \begin{pmatrix} C_{1,1} & C_{1,2} & C_{1,3} \\ C_{2,1} & C_{2,2} & C_{2,3} \\ C_{3,1} & C_{3,2} & C_{3,3} \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$$

for some 3×3 constant matrices $A = (A_{i,j}), B, C$ such that $\det A \cdot \det C \neq 0$, we will have the same system with new structure equations of the form (2) for $\tilde{R} = [\tilde{L}_1, \tilde{L}_2]$, $[\tilde{L}_j, \tilde{R}]$, $\tilde{R}^2$, but with transformed structure constants. We choose a continuous 1-parameter family of basis transformation matrices $A(\epsilon), B(\epsilon), C(\epsilon,\ 0 < \epsilon \leq 1$ such that $A(1) = C(1)$ is the identity matrix, $B(1) = 0$ and $\det A(\epsilon) \neq 0$, $\det C(\epsilon) \neq 0$. Now suppose as $\epsilon \to 0$ the basis change becomes singular, (i.e., the limits of A, B, C either do not exist or, if they exist do not satisfy $\det A(0) \det C(0) \neq 0$) but the structure equations involving $A(\epsilon), B(\epsilon), C(\epsilon)$, go to a limit, defining a new quadratic algebra Q'. We call Q' a *contraction* of Q in analogy with Lie algebra contractions [27].

For a degenerate superintegrable system with generators H, X, L_1, L_2 and structure equations (3),(4), defining a quadratic algebra Q, a change of basis to new generators $\tilde{H}, \tilde{X}\tilde{L}_1, \tilde{L}_2$ and parameter $\tilde{a}$ such that $\tilde{a} = Ca$, and

$$\begin{pmatrix} \tilde{L}_1 \\ \tilde{L}_2 \\ \tilde{H} \\ \tilde{X} \end{pmatrix} = \begin{pmatrix} A_{1,1} & A_{1,2} & A_{1,3} & 0 \\ A_{2,1} & A_{2,2} & A_{2,3} & 0 \\ 0 & 0 & A_{3,3} & 0 \\ 0 & 0 & 0 & A_{4,4} \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \\ X \end{pmatrix} + \begin{pmatrix} B_1 \\ B_2 \\ B_3 \\ 0 \end{pmatrix} a$$

for some 4×4 matrix $A = (A_{i,j})$ with $\det A \neq 0$, complex 4-vector B and constant $C \neq 0$ yields the same superintegrable system with new structure equations of the form (3),(4) for $[\tilde{X}, \tilde{L}_j]$, $[\tilde{L}_1, \tilde{L}_2]$, and $\tilde{G} = 0$, but with transformed structure constants. Suppose we choose a continuous 1-parameter family of basis transformation matrices $A(\epsilon), B(\epsilon), C(\epsilon)$, $0 < \epsilon \leq 1$ such that $A(1)$ is the identity matrix, $B(1) = 0$, $C(1) = 1$, and $\det A(\epsilon) \neq 0$, $C(\epsilon) \neq 0$. Now suppose as $\epsilon \to 0$ the basis change becomes singular, (i.e., the limits of

A, B, C either do not exist or, exist but do not satisfy $C(0) \det A(0) \neq 0$) but that the structure equations involving $A(\epsilon), B(\epsilon), C(\epsilon)$, go to a finite limit, thus defining a new quadratic algebra Q'. We call Q' a *contraction* of Q.

It has been established that all 2nd order 2D superintegrable systems can be obtained from system $S9$ by limiting processes in the coordinates and/or a Stäckel transformation, e.g. [30],[11] (2007a). All systems listed in Section 2 are limits of $S9$. It follows that the quadratic algebras generated by each system are contractions of the algebra of $S9$. (However, an abstract quadratic algebra may not be associated with a superintegrable system, and a contraction of a quadratic algebra associated with one superintegrable system to a quadratic algebra associated with another superintegrable system does not necessarily imply that this is associated with a coordinate limit process.)

4 Models of superintegrable systems

A representation of a quadratic algebra is a homomorphism of the algebra into the associative algebra of linear operators on some vector space. In this paper a *model* is a faithful representation in which the vector space is a space of polynomials in one complex variable and the action is via differential/difference operators acting on that space. We will study classes of irreducible representations realized by these models. Suppose a superintegrable system with quadratic algebra Q contracts to a superintegrable system with quadratic algebra Q' via a continuous family of transformations indexed by the parameter ϵ. If we have a model of an irreducible representation of Q we can try to "save" this representation by passing through a continuous family of irreducible representations of $Q(\epsilon)$ in the model to obtain a representation of Q' in the limit. We will show that as a byproduct of contractions to systems from $S9$ for which we save representations in the limit, we obtain the Askey Scheme for hypergeometric orthogonal polynomials. In all the models to follow the polynomials we classify are eigenfunctions of formally self-adjoint or formally skew-adjoint operators. To present compact results we will not derive the weight functions for the orthogonality. Most of them can be found in [28]. They can be found by requiring that the 2nd order operators H, L_1, L_2 are formally self-adjoint and the 1st order operator X is formally skew-adjoint. See [24] (2011) for some examples of this approach.

4.1 The $S9$ model

There is no differential model for $S9$ but a difference operator model yielding structure equations for the Racah and Wilson polynomials [24] (2007).

Recall, the Wilson polynomials are defined as

$$\begin{aligned} w_n(t^2) \equiv w_n(t^2,\alpha,\beta,\gamma,\delta) = (\alpha+\beta)_n(\alpha+\gamma)_n(\alpha+\delta)_n \times \qquad (26) \\ {}_4F_3\left(\begin{matrix} -n, & a+b+c+d+n-1, & a-t, & a+t \\ a+b, & a+c, & a+d & \end{matrix} ;1 \right) \\ = (a+b)_n(a+c)_n(a+d)_n \Phi_n^{(a,b,c,d)}(t^2), \end{aligned}$$

where $(a)_n$ is the Pochhammer symbol and ${}_4F_3(1)$ is a hypergeometric function of unit argument. The polynomial $w_n(t^2)$ is symmetric in $\alpha,\beta,\gamma,\delta$. For the finite dimensional representations the spectrum of t^2 is $\{(a+k)^2,\ k=0,1,\cdots,m\}$ and the orthogonal basis eigenfunctions are Racah polynomials. In the infinite dimensional case they are Wilson polynomials. They are eigenfunctions for the difference operator $\tau^*\tau$ defined via

$$\begin{aligned} \tau &= \frac{1}{2t}(E_t^{1/2}-E_t^{-1/2}), \\ \tau^* &= \frac{1}{2t}\left[(a+t)(b+t)(c+t)(d+t)E_t^{1/2}-(a-t)(b-t)(c-t)(d-t)E_t^{-1/2}\right], \end{aligned}$$

with $E_t^A F(t) = F(t+A)$.

A finite or infinite dimensional bounded below representation is defined by the following operators

$$L_1 = -4\tau^*\tau - 2(\alpha_2+1)(\alpha_3+1)+\frac{1}{2} \qquad (27)$$

$$L_2 = -4t^2+\alpha_1^2+\alpha_3^2-\frac{1}{2} \qquad (28)$$

$$H = E, \qquad (29)$$

where $a_i = \frac{1}{4}-\alpha_i^2$. The energy of the system is

$$E = -4(m+1)(m+1+\alpha_1+\alpha_2+\alpha_3)+2(\alpha_1\alpha_2+\alpha_1\alpha_3+\alpha_2\alpha_3)+\alpha_1^2+\alpha_2^2+\alpha_3^2-\frac{1}{4},$$

and the constants of the Wilson polynomials are chosen as

$$\begin{matrix} a = -\frac{1}{2}(\alpha_1+\alpha_3+1)-m, & d = \alpha_2+m+1+\frac{1}{2}(\alpha_1+\alpha_3+1), \\ b = \frac{1}{2}(\alpha_1+\alpha_3+1), & c = \frac{1}{2}(-\alpha_1+\alpha_3+1). \end{matrix}$$

Here $n = 0,1,\cdots,m$ if m is a nonnegative integer and $n = 0,1,\cdots$ otherwise.

Taking a basis as

$$f_{n,m} \equiv \Phi_n^{(a,b,c,d)}(t^2),$$

the action of the model on the basis is

$$\begin{aligned}
L_1 f_{n,m} &= -\left(4n^2+4n[\alpha_2+\alpha_3+1]+2[\alpha_2+1][\alpha_3+1]-\frac{1}{2}\right) f_{n,m} \qquad (30)\\
L_2 f_{n,m} &= K(n+1,n)f_{n+1,m}+K(n-1,n)f_{n-1,m}+\left(K(n,n)+\alpha_1^2+\alpha_3^2-\frac{1}{2}\right) f_{n,m}\\
H f_{m,n} &= E f_{n,m}.
\end{aligned}$$

with

$$K(n+1,n)=\frac{(\alpha_3+1+\alpha_2+n)\,(m-n)\,(m-n+\alpha_1)\,(1+\alpha_2+n)}{(\alpha_3+1+\alpha_2+2\,n)\,(\alpha_3+2+\alpha_2+2\,n)},$$

$$K(n-1,n)=\frac{n\,(\alpha_3+n)\,(\alpha_1+\alpha_3+1+\alpha_2+m+n)\,(1+\alpha_3+\alpha_2+m+n)}{(\alpha_3+1+\alpha_2+2\,n)\,(\alpha_3+\alpha_2+2\,n)},$$

$$K(n,n)=\left(\frac{1}{2}(\alpha_1+\alpha_3+1)-m\right)^2-K(n+1,n)-K(n-1,n).$$

5 Nondegenerate to nondegenerate limits

5.1 Contractions $S9 \to E1$

There are at least two ways to take this contraction; it is possible to contract the sphere about the point $(0,1,0)$ which gives the contraction of representation in terms of Wilson polynomials to continuous Dual Hahn polynomials. Contracting about the point $(1,0,0)$ leads to continuous Hahn polynomials or Jacobi polynomials. The dual Hahn and continuous dual Hahn polynomials correspond to the same superintegrable system but they are eigenfunctions of different generators. For the finite dimensional restrictions (m a positive integer) we have the restrictions of Racah polynomials to Dual Hahn and Hahn respectively.

1) Wilson → Continuous Dual Hahn For the first limit, in the quantum system, we contract about the point $(0,1,0)$ so that the points of our two dimensional space lie in the plane $(x,1,y)$. We set $s_1=\sqrt{\epsilon}x, s_2=\sqrt{1-s_1^2-s_3^2}\approx 1-\frac{\sqrt{\epsilon}}{2}(x^2+y^2)$, $s_3=\sqrt{\epsilon}y$, for small ϵ. The coupling constants are transformed as

$$\begin{pmatrix}\tilde a_1\\ \tilde a_2\\ \tilde a_3\end{pmatrix}=\begin{pmatrix}0&\epsilon^2&0\\1&0&0\\0&0&1\end{pmatrix}\begin{pmatrix}a_1\\ a_2\\ a_3\end{pmatrix}, \qquad (31)$$

and we get $E1$ as $\epsilon \to 0$. This gives the quadratic algebra contraction defined by the contraction of the operators

$$\begin{pmatrix} \tilde{L}_1 \\ \tilde{L}_2 \\ \tilde{H} \end{pmatrix} = \begin{pmatrix} \epsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \epsilon \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -\epsilon & 0 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}. \tag{32}$$

As in S9, it is advantageous in the model to express the 3 coupling constants as quadratic functions of other parameters, so that

$$\begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix} = \begin{pmatrix} -\beta_1^2 \\ \frac{1}{4} - \beta_2^2 \\ \frac{1}{4} - \beta_3^2 \end{pmatrix} = \begin{pmatrix} \frac{\epsilon^2}{4} - \epsilon^2 \alpha_2^2 \\ \frac{1}{4} - \alpha_1^2 \\ \frac{1}{4} - \alpha_3^2 \end{pmatrix} \tag{33}$$

with $\alpha_2 \to \infty$.

In the contraction limit the operators tend to

$$L_1' = \lim_{\epsilon \to 0} \widetilde{L}_1 = -4\tau'^{*}\tau' - 2\beta_1(\beta_3 + 1) \tag{34}$$

$$L_2' = \lim_{\epsilon \to 0} \widetilde{L}_2 = -4t^2 + \beta_2^2 + \beta_3^2 - \frac{1}{2} \tag{35}$$

$$H' = \lim_{\epsilon \to 0} \widetilde{H} = E'. \tag{36}$$

The energy of the system is now

$$E' = -2\,\beta_1\,(2\,m + 2 + \beta_2 + \beta_3)\,.$$

The eigenfunction of L_1, the Wilson polynomials, transform in the contraction limit to the eigenfunctions of L_1', which are the dual Hahn polynomials S_n,

$$S_n(-t^2, a', b', c') = (a' + b')_n (a' + c')_n {}_3F_2 \left(\begin{matrix} -n, & a' + t, & a' - t \\ a' + b', & a' + c' & \end{matrix} ; 1 \right)$$

where the constants of the dual Hahn polynomials are

$$a' = -\frac{1}{2}(\beta_2 + \beta_3 + 1) - m, \quad b' = \frac{1}{2}(\beta_2 + \beta_3 + 1), \quad c' = \frac{1}{2}(-\beta_2 + \beta_3 + 1).$$

Again, $n = 0, 1, \cdots, m$ if m is a nonnegative integer and $n = 0, 1, \cdots$ otherwise. The operators τ'^{*} and τ' are given by

$$\tau' = \tau = \frac{1}{2t}(E_t^{1/2} - E_t^{-1/2}),$$

$$\tau^{*} = \frac{\beta_1}{2t}\left[(a' + t)(b' + t)(c' + t)E_t^{1/2} - (a' - t)(b' - t)(c' - t)E_t^{-1/2}\right],$$

Taking a basis as

$$f'_{n,m} \equiv \frac{S_n(-t^2, a', b', c')}{(a'+b')_n (a'+c')_n},$$

the action of the model on the basis is

$$\begin{aligned} L'_1 f'_{n,m} &= -2\beta_1 \left(2n+\beta_3+1\right) f'_{n,m} \\ L'_2 f'_{n,m} &= K'(n+1,n) f'_{n+1,m} + K'(n-1,n) f'_{n-1,m} + \left(K'(n,n)+\beta_2^2+\beta_3^2-\frac{1}{2}\right) f_{n,m} \\ H' f_{m,n} &= E' f_{n,m}. \end{aligned}$$

with

$$K'(n+1,n) = (m-n)\left(m-n+\beta_2\right), \qquad K'(n-1,n) = n(n+\beta_3)$$

$$K'(n,n) = \left(\frac{1}{2}(\beta_2+\beta_3+1)-m\right)^2 - K'(n+1,n) - K'(n-1,n).$$

2) Wilson → Continuous Hahn For the next limit, we contract about the point $(1,0,0)$ so that the points of our two dimensional space lie in the plane $(1,x,y)$. We set $s_1 = \sqrt{1-s_1^2-s_3^2} \approx 1 - \frac{\sqrt{\epsilon}}{2}(x^2+y^2)$, $s_2 = \sqrt{\epsilon}x$, $s_3 = \sqrt{\epsilon}y$, for small ϵ. The coupling constants are transformed as

$$\begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix} = \begin{pmatrix} \epsilon^2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, \tag{37}$$

and we get $E1$ as $\epsilon \to 0$. This gives the quadratic algebra contraction

$$\begin{pmatrix} \tilde{L}_1 \\ \tilde{L}_2 \\ \tilde{H} \end{pmatrix} = \begin{pmatrix} 0 & \epsilon & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \epsilon \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -\epsilon & 0 & 0 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}. \tag{38}$$

In terms of the constants of (33) the transformation gives

$$\begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix} = \begin{pmatrix} -\beta_1^2 \\ \frac{1}{4}-\beta_2^2 \\ \frac{1}{4}-\beta_3^2 \end{pmatrix} = \begin{pmatrix} \frac{\epsilon^2}{4}-\epsilon^2\alpha_1^2 \\ \frac{1}{4}-\alpha_2^2 \\ \frac{1}{4}-\alpha_3^2 \end{pmatrix}, \tag{39}$$

with $\alpha_1 \to \infty$.

Saving a representation: We set

$$t = -x + \frac{\beta_1}{2\epsilon} + m + \frac{1}{2}(\beta_3+1). \tag{40}$$

In the contraction limit, the operators go as $L'_i = \lim_{\epsilon\to 0} \widetilde{L}_i$,

$$\begin{aligned}
L'_1 &= 2\beta_1\,(2\,x - 2m - \beta_3 - 1) \\
L'_2 &= -4\left(B(x)E_x + C(x)E_x^{-1} - B(x) - C(x)\right) - 2(\beta_2+1)(\beta_3+1) + \frac{1}{2} \\
H' &= -2\beta_1(2m + 2 + \beta_2 + \beta_3). \qquad (41)
\end{aligned}$$

with $B(x) = (x-m)(x+\beta_2+1)$, $C(x) = x(x-m-1-\beta_3)$. The operators L'_1, L'_2 and H' satisfy the algebra relations in (8). The eigenfunction of L_1, the Wilson polynomials, transform in the contraction limit to the eigenfunctions of L'_2, which are the Hahn polynomials, $f'_{m,n} = Q_n$,

$$Q_n(x;\beta_2,\beta_3,m) = {}_3F_2\left(\begin{matrix} -n, & \beta_2+\beta_3+n+1, & -x \\ -m, & \beta_2+1 & \end{matrix}\; ;1\right).$$

The action of the operators on this basis is given by

$$\begin{aligned}
L'_1 f'_{n,m} &= K'(n+1,n)f'_{n+1,m} + K'(n,n)f'_{n,m} + K'(n-1,n)f'_{n-1,n}, \\
L'_2 f'_{n,m} &= -\left(4n^2 + 4n[\beta_2+\beta_3+1] + 2[\beta_2+1][\beta_3+1] - \frac{1}{2}\right) f'_{n,m}.
\end{aligned}$$

$$\begin{aligned}
K'(n+1,n) &= -4\beta_1\frac{(m-n)(n+\beta_2+\beta_3+1)(n+\beta_2+1)}{(2n+\beta_2+\beta_3+1)(2n+\beta_2+\beta_3+2)} \\
K'(n-1,n) &= -4\beta_1\frac{n(n+\beta_3)(m+n+\beta_2+\beta_3+1)}{(2n+\beta_2+\beta_3+1)(2n+\beta_2+\beta_3)} \\
K'(n,n) &= -2\beta_1(2m+\beta_3+1) - K'(n+1,n) - K(n-1,n).
\end{aligned}$$

If $\beta_2 = \bar{\beta}_3$ there is still a real 3-term recurrence relation. In the original system the potential is PT-symmetric even though complex, so the energy spectrum is real. In this case one studies the original system and its dual and obtains biorthogonality, rather than an orthonormal basis.

3) Wilson $\to$ Jacobi: The previous contraction on the operator realization degenerates when $\alpha_1 = \beta_1 = 0$. However, we can save this representation by setting

$$m = \frac{\sqrt{-E'}}{2\sqrt{\epsilon}} - 1 + \frac{\beta_2+\beta_3}{2}, \qquad t = \frac{\sqrt{-E'}}{2\sqrt{\epsilon}}\sqrt{\frac{1+x}{2}}$$

for E' a constant and letting $m \to \infty$. Then, the contraction (38) gives a contraction of the model for S9 to a differential operator model for $E1$ with $\beta_1 = 0$:

$$\begin{aligned}
H' &= E', \\
L'_1 &= \tfrac{E'}{2}(x+1), \qquad (42) \\
L'_2 &= 4(1-x^2)\partial_x^2 + 4\left[\beta_3 - \beta_2 - (\beta_2+\beta_3+2)x\right]\partial_x - 2(\beta_2+1)(\beta_3+1) + \tfrac{1}{2}.
\end{aligned}$$

The eigenfunctions for L_1, the Wilson polynomials, tend in the limit to eigenfunction of L_2' which are the Jacobi polynomials:

$$P_n^{\beta_2,\beta_3}(x) = \frac{(\beta_2+1)_n}{n!} {}_2F_1\left(\begin{array}{cc} -n, & \beta_2+\beta_3+n+1 \\ \beta_2+1 & \end{array}; \frac{x-1}{2}\right). \qquad (43)$$

Taking a basis as $f_n = \frac{n!}{(\beta_2+1)_n} P_n^{\beta_2,\beta_3}(x)$ the action of the operators on the basis is

$$\begin{aligned} L_1' f_n' &= K'(n+1,n) f_{n+1}' + K'(n-1,n) f_{n-1} + K'(n,n) f_n \\ L_2' f_n' &= -4n(n+\beta_2+\beta_3+1) - 2(\beta_2+1)(\beta_3+1) + \frac{1}{2}, \end{aligned}$$

with

$$\begin{aligned} K'(n+1,n) &= \frac{E'(\beta_2+\beta_3+n+1)(\beta_2+n+1)}{(\beta_2+\beta_3+2n+1)(\beta_2+\beta_3+2n+2)}, \\ K'(n-1,n) &= \frac{E'n(n+\beta_3)}{(\beta_2+\beta_3+2n)(\beta_2+\beta_3+2n+1)}, \qquad (44) \\ K'(n,n) &= E' - K'(n+1,n) - K'(n-1,n). \qquad (45) \end{aligned}$$

If $\beta_2 = \bar{\beta}_3$ then there is a real 3-term recurrence relation and the potential is PT-symmetric though complex. In this case the basis functions above, suitably renormalized, are pseudo Jacobi polynomials [29]. If $\beta_2 = \bar{\beta}_3$ there is a real 3-term recurrence relation and the potential is PT-symmetric so E is real. In this case one studies the original system and its dual and obtains biorthogonality of the basis functions.

5.2 Contractions $E1 \to E8$

1) Hahn → Jacobi: Here, we express 2D Euclidean space in complex variables and let $z \to \infty$, $\overline{z} \to 0$ as

$$x = \frac{1}{2}\left(\sqrt{\epsilon} z + \overline{z}\epsilon^{-\frac{1}{2}}\right), \qquad y = \frac{-i}{2}\left(\sqrt{\epsilon} z - \overline{z}\epsilon^{-\frac{1}{2}}\right).$$

The coupling constants are transformed as

$$\begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -8\epsilon^2 & -8\epsilon^2 \\ 0 & 4\epsilon & -4\epsilon \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, \qquad (46)$$

and the system E8 is obtained by the following singular limit:

$$\begin{pmatrix} \tilde{L}_1 \\ \tilde{L}_2 \\ \tilde{H} \end{pmatrix} = \begin{pmatrix} \epsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}. \tag{47}$$

Consider the second $E1$ model above, based on the Hahn polynomials. For simplicity of the model, we introduce new parameters γ_i as in

$$\begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix} = \begin{pmatrix} -\gamma_1^2 \\ 16\gamma_3^2 + \mathcal{O}(\epsilon) \\ 8\gamma_3(\gamma_2 - 2) + \mathcal{O}(\epsilon) \end{pmatrix} = \begin{pmatrix} -\beta_1^2 \\ -4\epsilon^2(1 - 2\beta_2^2 - 2\beta_3^2) \\ -4\epsilon(\beta_2^2 - \beta_3^2) \end{pmatrix}. \tag{48}$$

We save the representation via (48) and the following change of variable

$$x = \frac{\gamma_3}{\epsilon}\left(\frac{1-t}{2}\right).$$

In the contraction limit $L_i' = \lim_{\epsilon\to 0} \tilde{L}_i$ (47), the model becomes

$$\begin{aligned} L_1' &= -2\gamma_1\gamma_3 t \\ L_2' &= 4(1-t^2)\partial_t^2 + 4(2m + \gamma_2 - \gamma_2 t)\partial_t - (\gamma_2 - 1)^2 \\ H' &= -2\gamma_1(2m + \gamma_2). \end{aligned} \tag{49}$$

The eigenfunctions for L_2 (41), Hahn polynomials, tend in the limit to eigenfunctions of L_2', Jacobi polynomials

$$P_n^{-m-1,\gamma_2+m-1}(t) = \frac{(-m)_n}{n!}\,{}_2F_1\left(\begin{matrix} -n & n+\gamma_2+1 \\ & -m \end{matrix} \,\Big|\, \frac{1-t}{2}\right),$$

with the normalization $\lim_{\epsilon\to 0} f_{m,n} = f_{m,n}' = \frac{n!}{(-m)_n} P_n^{-m-1,\gamma_2+m-1}(t)$. The action of the operators on this basis becomes

$$\begin{aligned} L_1' f_n' &= K'(n+1,n) f_{n+1}' + K'(n-1,n) f_{n-1} + K'(n,n) f_n \\ L_2' f_n' &= \left[-4n(n+\gamma_2-1) - (\gamma_2-1)^2\right] f_n', \end{aligned}$$

with

$$\begin{aligned} K'(n+1,n) &= -\frac{4\gamma_1\gamma_3(m-n)(n+\gamma_2-1)}{(2n+\gamma_2-1)(2n+\gamma_2)} \\ K'(n-1,n) &= \frac{4\gamma_1\gamma_3 n(n+m+\gamma_2-1)}{(2n+\gamma_2-1)(2n+\gamma_2-2)} \\ K(n,n) &= -2\gamma_1\gamma_3 - K(n+1,n) - K(n-1,n). \end{aligned}$$

Note that this model gives a finite dimensional representation in the case that m is a positive integer. This is in contradistinction to the previous model based on Jacobi polynomials (43) which gives only infinite dimensional representations. This is in agreement with the fact that the physical system E1 with $a_1 = 0$ has only unbounded trajectories.

2) Jacobi → generalized Bessel polynomials: The same contraction (47) acting on the model for E1 with $a_1 = 0$ (42), gives a model based on the generalized Bessel polynomials [32]:

$$\begin{aligned} L_1' &= -\gamma_3 E' t \\ L_2' &= -4t^2\partial_t^2 - 4(1+\gamma_2 t)\partial_t - (\gamma_2 - 1)^2 \\ H' &= E', \end{aligned} \tag{50}$$

where we have used the change of variable $x = -2\gamma_3 t/\epsilon$.

5.3 Contraction $E1 \to E3'$

1) Dual Hahn → Meixner, Krawtchouk, and Meixner-Pollaczek: In the quantum $E1$ system we make a change of variables

$$x \to x - \frac{8(-c)^{\frac{1}{4}}a_1 - a_2}{8\epsilon a_1} - \frac{c}{8\epsilon^5}, \qquad y \to y - \frac{8a_1(-c)^{-\frac{1}{4}} - a_3}{8\epsilon a_1} - \frac{b}{8c\epsilon^5}.$$

To cancel the singularities, the constants for the new system are chosen as

$$\begin{pmatrix} \tilde a_1 \\ \tilde a_2 \\ \tilde a_3 \end{pmatrix} = \begin{pmatrix} 4 & 0 & 0 \\ c\epsilon^{-5} & \epsilon^{-1} & 0 \\ c\epsilon^{-5} & 0 & \epsilon^{-1} \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, \tag{51}$$

and the new operators are

$$\begin{pmatrix} \tilde L_1 \\ \tilde L_2 \\ \tilde H \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -(-c)^{-\frac{1}{2}}(c+1) & \epsilon^2 & (-c)^{-\frac{1}{2}} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \end{pmatrix} + \begin{pmatrix} -2(-c)^{-\frac{1}{2}}\epsilon^{-2} & 0 & 0 \\ -2\epsilon^{-2} & 0 & 0 \\ 2\epsilon^{-2}(-c)^{-\frac{1}{2}}(c-1) & 0 & 0 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}. \tag{52}$$

In the limit as $\epsilon \to 0$ the operators and algebra tend to those of E3' (12). Note that the constant c is not inherent in either model but instead parameterizes a family of contractions and leads to a parameterized family of models.

In the quantum $E1$ system we let $x \to x + \sqrt{\frac{c_2}{\epsilon\omega}} - \frac{b}{8\omega^2}$, $y \to y + \sqrt{\frac{c_1}{\epsilon\omega}} - \frac{c}{8\omega^2}$, and go to the contraction limit. We find $L_1' = \partial_x^2 - 4\omega^2 x^2 + bx - \frac{b^2}{16\omega^2} + 2\omega$, $L_2' = \partial_y^2 - 4\omega^2 y^2 + cy - \frac{c^2}{16\omega^2} + 2\omega$, $L_3' = \frac{c_1}{\omega} L_1' + \frac{c_2}{\omega} L_2' - 2\frac{\sqrt{c_1 c_2}}{w}\left(\partial_x\partial_y - 4\omega^2 xy + \frac{bx+cy}{2} - \frac{bc}{16\omega^2}\right)$. Up to a change of basis, this is exactly the system $E3'$. Note that ω is rescaled by the contraction.

At the quadratic algebra level it is convenient to take a non=canoncal version of the contraction: Consider the first $E1$ model above. Set $B_1 = c_1/\epsilon$, $B_2 = c_2/\epsilon$, and $L_3' = \epsilon\left(L_3 + 2(\frac{c_1}{\epsilon}+1)(\frac{c_2}{\epsilon}+1)\right)$, $L_1' = L_1 + 2\omega(\frac{c_2}{\epsilon}+1)$, $L_2' = L_2 + 2\omega(\frac{c_1}{\epsilon}+1)$, $H' = L_1' + L_2'$, $\epsilon R' = [L_1', L_3'] = \epsilon[L_1, L_3]$. In the contraction limit we have

$$f_{n,m} = {}_2F_1\left(\begin{matrix} -n, & -x \\ -m, & \end{matrix} ; 1 - \frac{1}{c}\right), \tag{53}$$

the Meixner polynomials as limits of Hahn polynomials. Here, $c = -c_2/c_1$. Further,

$$L_2' f_{n,m} = 4x\omega f_{n,m} = -4K(n+1,n) f_{n+1,m} - 4K(n,n) f_{n,m} - 4K(n-1,n) f_{n-1,n},$$

$$K(n+1,n) = \frac{\omega(m-n)c_2}{c_1+c_2},\quad K'(n-1,n) = \frac{\omega n c_1}{c_1+c_2},\quad H' = -4\omega m.$$

$$L_1' f_{n,m} = 4n(c_1+c_2) f_{n,m} = -4\left(B(x)T_x + C(x)T_x^{-1} - [B(x)+C(x)]\right) f_{n,m},$$

$$B(x) = c_2(x-m),\quad C(x) = -c_1 x,\quad K'(n,n) = -K'(n+1,n) - K'(n-1,n).$$

In the case where m is a positive integer the Meixner polynomials reduce to Krawtchouk polynomials.

Recall that in the model for the system E1, the dual Hahn polynomials had a real 3-term recurrence relation when the system itself was PT-symmetric. If we retain this restriction in the limit, the constant c is required to have modulus 1, $c = e^{2i\phi}$, and $a_2' = \overline{a_3'}$ in the physical system. In this case, the Meixner-Pollaczek polynomials are obtained as a limit of the dual Hahn

$$P_n^{\lambda}(x;\phi) = \frac{(-m)_n}{n!} e^{in\phi}{}_2F_1\left(\begin{matrix} -n, & -\frac{m}{2} + ix \\ -m, & \end{matrix} ; 1 - e^{-2i\phi}\right).$$

Here, we have made a change of variables $x \to ix + m/2$.

The related $E3'$ quantum system has special properties. We choose it as

$$H = \partial_{xx} + \partial_{yy} - \omega^2(x^2+y^2) + a_2' x + \overline{a_2}' y - \frac{a_2'^2 + \overline{a_2'}^2}{4\omega^2}$$

where $\omega > 0$. This system admits PT-symmetry; the potential V is complex but the bound-state eigenvalues are real:

$$E_m = -2\omega(n_1 + n_2 + 1), \qquad n_1 + n_2 = m = 0, 1, 2, \cdots .$$

Here, H is not self-adjoint but its basis vectors and the basis vectors of its adjoint H^* are biorthogonal.

2) Hahn $\to$ Meixner, Krawtchouk, and Meixner-Pollaczek: We use the same limit as immediately above, but apply it to the second $E1$ model (41) to obtain

$$\begin{aligned} H' &= -2\omega(m+1) \qquad (54)\\ L_1' &= -\omega(2x-2m-1)\\ L_2' &= \frac{2\omega}{\sqrt{-c}}\left[(x-m)T_x^1 + 2T_x^{-1}\right]. \end{aligned}$$

The operator which is diagonalized by the Meixner polynomials (53) is $c\sqrt{-c}L_2' + (1+c)L_1 - cH'$. The above discussion of the PT-symmetric limit also applies for this model.

3) Krawtchouk $\to$ Charlier: Beginning with the model based on the Krawtchouk basis (53) and consider the $m \to \infty$ limit. This is a different type of contraction than considered above because the $m+1$ dimensional eigenspace is changing with each increment of m. We can save the representation by taking $c_2 = 1/m$, $c_1 = -(1/a + 1/m)$. The basis functions become

$$f_n(x) = C_n(x;a) = \lim_{m\to\infty} {}_2F_1\left(\begin{matrix} -n, & -x \\ m & \end{matrix}; -\frac{m}{a}\right) = {}_2F_0\left(\begin{matrix} -n, & -x \\ - & \end{matrix}; -\frac{1}{a}\right),$$

the Charlier polynomials. The difference operator determining these polynomials and the three term recurrence relation are obtained in the limit:

$$\hat{L}_2' f_n = (L_2' + 4\omega m)f_n = 4x\omega f_n = -4K(n+1,n)f_{n+1} - 4K(n,n)f_n - 4K(n-1,n)f_{n-1},$$

$$K(n+1,n) = -\omega a, \quad K(n-1,n) = \omega n, \quad K(n,n) = \omega(a-n),$$

$$L_3 f_n = -\frac{4n}{a}f_n = -4\left(B(x)T_x + C(x)T_x^{-1} - [B(x)+C(x)]\right)f_n,$$

$$B(x) = -1, \quad C(x) = \frac{x}{a}, \ \hat{H}' = H' + 4\omega m = 0.$$

As we go to the contraction limit the model is restricted to the eigenspace of H' with eigenvalue $-4\omega m$, i.e., on the eigenspace of $\hat{H}'$ with eigenvalue 0. In the quantum system, the Hamiltonian $H' + 4\omega m$ and the symmetry $L_2' + 4\omega m$ blow up with m, so don't give a finite limit. However, the quadratic algebra converges in this contraction of E3' to itself. The model is giving us asymptotic information in m about the relation between $L2'$ and $L1'$ eigenbases of H' on the $-4\omega m$ eigenspace.

5.4 Contraction $E1 \to E2$:

In quantum $E1$ we let $x \to x + \sqrt{\frac{c_2}{\omega\epsilon}} - \frac{b}{8\omega^2}$, $y \to y$, and go to the limit. This induces a algebra contraction to $E2$: Let $B_1 = c_2/\epsilon$ and $L_1' = L_1 + 2\frac{\omega c_2}{\epsilon} + \frac{b^2}{16\omega^2}$, $L_2' = L_2$, $L_3' = \frac{1}{2}\sqrt{\frac{\omega\epsilon}{c_2}} L_3 + \frac{1}{8\omega^2} L_2 - \frac{1}{2}\sqrt{\frac{c_2}{\omega\epsilon}} L_2$, $H' = L_1' + L_2' = H + \frac{2\omega c_2}{\epsilon} + \frac{b^2}{16\omega^2}$, $R' = \sqrt{\epsilon} R$. We can't save the representation.

5.5 Contraction $E8 \to E10$

In the physical model we translate to infinity: $z = z' - 1/\epsilon^2$, $\bar{z} = \bar{z}' + 1/\epsilon$ and cancel the singularities that occur. The coupling constants transform as

$$\begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix} = \begin{pmatrix} \frac{-4a_1}{3} & 0 & -\frac{8a_3\epsilon^5}{3} \\ \frac{4a_1}{3\epsilon} & 0 & \frac{2a_3\epsilon^4}{3} \\ \frac{-a_2}{\epsilon^2} & 3\epsilon^2 a_2 & -2\epsilon^3 a_3 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, \tag{55}$$

and the system E10 is obtained by the following singular limit:

$$\begin{pmatrix} \tilde{L}_1 \\ \tilde{L}_2 \\ \tilde{H} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -\frac{1}{\epsilon^2} & \epsilon^2 & \frac{1}{2\epsilon} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \end{pmatrix} + \begin{pmatrix} \frac{1}{6\epsilon^2} & 0 & -\frac{\epsilon^3}{6} \\ \frac{1}{4\epsilon^4} & -\frac{1}{4} & \frac{\epsilon}{2} \\ \frac{1}{\epsilon^3} & \epsilon & -\epsilon^2 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}. \tag{56}$$

We can not save the representation.

6 Nondegenerate to degenerate contractions

This appears initially a mere restriction of the 3-parameter potential to 1-parameter. However, after restriction one 2nd order generator L_i becomes a perfect square $L_i = X^2$. The spectrum of L_i is nonnegative but that of X can take both positive/negative values. This results in a virtual doubling of the support of the measure in the finite case. Also, the commutator of X and the remaining 2nd order symmetry leads to a new 2nd order symmetry. In [26] we will show how this Casimir follows directly as a contraction from the expression for R^2.

6.1 Contraction $S9 \to S3$:

1) Wilson → special dual Hahn (1st model): The quantum system $E3$ (2.1) is given in the singular limit from system $S9$ (26) by

$$a_2 = a_3 = \epsilon \to 0, \qquad a_1 = a_1 \tag{57}$$

$$X'^2 = \lim_{\epsilon\to 0} L_1, \qquad L_2' = \lim_{\epsilon\to 0} L_2, \qquad L_1' = [X', L_2']. \tag{58}$$

The operators in this contraction differ from those given in subsection 2.1 by a cyclic permutation of the coordinates $s_i \to s_{i+1}$.

Now we investigate how the difference operator realization of $S9$ contracts to irreducible representations of the $S3$ symmetry algebra. This is more complicated since the original restricted algebra is now contained as a proper subalgebra of the contracted algebra.

The contraction (71) is realized in the model by setting $\alpha_2 = \alpha_3 = -1/2$ and $\alpha_1 = \alpha$ (the subscript is dropped in this model since there is now a sole α). The restricted operators then become $H' = E'$ with $E' = -4(m+1)(m+\alpha) - (\alpha-1)^2 + \frac{1}{4}$ and

$$X'^2 = -4\tau^*\tau, \qquad L'_2 = -4t^2 + \alpha^2 - \frac{1}{4}. \tag{59}$$

The eigenfunctions for X'^2, the Wilson polynomials, become

$$\Phi_{\pm n}(t^2) = {}_4F_3\left(\begin{array}{llll} -n, & n, & -\frac{4m+2\alpha+1}{4} - t, & -\frac{4m+2\alpha+1}{4} + t \\ -m, & -m-\alpha, & \frac{1}{2} & \end{array} ;1 \right). \tag{60}$$

Here $n = 0, 1, \cdots, m$ if m is a nonnegative integer and $n = 0, 1, \cdots$ otherwise. For finite dimensional representations, the spectrum of t is the set $\{\frac{\alpha}{2} + \frac{1}{4} + m - k,\ k = 0, 1, \cdots, m\}$. Note that the restricted polynomial functions (60) are no longer the correct basis functions for the contracted superintegrable system. To see this, we consider the contracted expansion coefficients

$$L'_2 f_{n,m} = K(n+1,n) f_{n+1,m} + K(n-1,n) f_{n-1,m} + \left(K(n,n) + \alpha_1^2 - \frac{1}{4}\right) f_{n,m}$$

with

$$\begin{aligned} K(n+1,n) &= \frac{1}{4}(m-n+\alpha)(m-n) && (61)\\ K(n-1,n) &= \frac{1}{4}(m+n+\alpha)(m+n) && (62)\\ K(n,n) &= \left(\frac{1}{2}(\alpha+\frac{1}{2}) - m\right)^2 - K(n+1,n) - K(n-1,n). && (63) \end{aligned}$$

Indeed $K(n-1,n)$ no longer vanishes for $n = 0$, so $f_{0,m}$ is no longer the lowest weight eigenfunction, note $f_{-1,m} = f_{1,m}$ is still a polynomial in t^2. To understand the contraction we set $n = N - M/2$ where N is a nonnegative integer and $M = 2m$. Then the equations for the K's become

$$\begin{aligned} K(N+1,N) &= \frac{1}{4}(M-N+\alpha)(M-N) && (64)\\ K(N-1,N) &= \frac{1}{4}N(N+\alpha) && (65) \end{aligned}$$

the three term recurrence relation gives a new set of basis orthogonal polynomials for representations of $S3$. The lowest eigenfunction occurs for $N=0$; if M is a nonnegative integer the representation is $2m+1$-dimensional with highest eigenfunction for $N=M$. The basis functions which satisfy this three-term recurrence are

$$f_{N,M}(t^2)=\frac{(\alpha+1)_N}{(-\alpha-M)_N}{}_3F_2\left(\begin{array}{ccc}-N, & -s, & s+2\alpha+1\\ -M, & 1+\alpha & \end{array};1\right). \tag{66}$$

Here f_N is a polynomial of order $2N$ in s and of order n in $\lambda(s)=s(s+2\alpha+1)$, a special case of dual Hahn polynomials. These dual Hahn polynomials satisfy the following difference operator,

$$X=i\left(B(s)E_s+C(s)E_s^{-1}\right), \tag{67}$$

with $B(s)+C(s)=M$ defined as

$$B(s)=\frac{(s+2\alpha+1)(M-s)}{2s+2\alpha+1}, \qquad C(s)=\frac{s(s+M+2\alpha+1)}{2s+2\alpha+1}.$$

The operators which form a model for the algebra (2.1) are X (67) along with

$$L_1=-\left(s+\alpha+\frac{1}{2}\right)^2+\alpha^2-\frac{1}{4}, \qquad L_2=[L_1,X]. \tag{68}$$

For finite dimensional representations the spectrum of s is $\{0,1,\cdots,M\}$. From the operator L_1 we can determine the relation between t and s, namely $s=2t-\alpha-1/2$.

What is the relation between the functions (60) and the proper basis functions (66)? Note this model X,L_1,L_2 can be obtained from the contracted model X',L_1',L_2' by conjugating by the "ground state" of the contracted model $\Phi_{-\frac{M}{2}}(t^2)$. We find explicitly the gauge function

$$\Phi_{-\frac{M}{2}}(t^2)=\frac{(\frac{1}{2}-\frac{M}{2})_k(-\alpha-M)_k}{(\frac{1}{2})_k(-\alpha-\frac{M}{2})_k},$$

when t is evaluated at the weights

$$t=\frac{\alpha}{2}+\frac{1}{4}+\frac{M}{2}-k, \quad k=0,1,\cdots\frac{M}{2}. \tag{69}$$

So, the operator X' is related to X via conjugation by $\Phi_{-\frac{M}{2}}(t^2)$.

Note that the functions $\Phi_n(t^2)$ are only defined for discrete values of t (69). However, on this restricted set the functions $\Phi_{-\frac{M}{2}+N}$ and $f_{N,M}$ satisfy exactly the same three term recurrence formula under multiplication by $-4t^2 - a$, with the bottom of the weight ladder at $N = 0$. From this we find the identity

$$\Phi_{-\frac{M}{2}}(t^2) f_{N,M}(t^2) = \Phi_{-\frac{M}{2}+N}(t^2), \qquad t = \frac{\alpha}{2} + \frac{1}{4} + \frac{M}{2} - k. \tag{70}$$

Since $\Phi_{-\frac{M}{2}}(t^2) = \Phi_{\frac{M}{2}}(t^2)$ this relation implies that $f_{M,M}(t^2) = 1$ when restricted to the spectrum of t.

2) Wilson → special Hahn (2nd model): The quantum system $S3$ (16) can also be obtained from system $S9$ (26) by

$$\begin{aligned} a_1 = a_3 = \epsilon \to 0, \qquad a_2 = \frac{1}{4} - \alpha^2 \\ L_1' = \lim_{\epsilon\to 0} L_1, \qquad X'^2 = \lim_{\epsilon\to 0} L_2, \qquad L_2' = [X', L_1']. \end{aligned} \tag{71}$$

Again, the physical model obtained by this contraction is related to the that given in subsection 2.1 by a cyclic permutation of the coordinates $s_i \to s_{i-1}$.

In this limit, the operator X^2 can be immediately factorized to obtain the skew-adjoint operator $X = 2it$. Taking $x = t + m$, the operators in our model are given by

$$\begin{aligned} L_1' = & \; -\left[B(x)E_x + C(x)E_x^{-1} - B(x) - C(x)\right] - \alpha - \tfrac{1}{2} \\ & \; B(x) = (x - 2m)(x + \alpha + 1), \qquad C(x) = x(x - 2m - \alpha - 1), \\ X' = & \; 2i(x - m), \qquad L_2' = [X', L_1] \end{aligned} \tag{72}$$

which is diagonalized by Hahn polynomials

$$\widehat{f}_{k,m} = {}_3F_2\left(\begin{matrix} -k, k + 2\alpha + 1, -x \\ \alpha + 1, -2m \end{matrix} ; 1 \right) = Q_k(x; B_2, B_2, 2m), \tag{73}$$

$$L_1' \widehat{f}_{k,m} = \left(-\left(k + \alpha + \frac{1}{2} \right)^2 + \alpha^2 - \frac{1}{4} \right) \widehat{f}_{k,m}, \quad k = 0, 1, \cdots, 2m. \tag{74}$$

These polynomials satisfy special recurrence relations not obeyed by general Hahn polynomials. Note that the dimension of the representation space has jumped from $m+1$ to $2m+1$. Comparing these eigenfunctions with the limit of the Wilson polynomials,

$$\lim_{\epsilon\to 0} L_1' f_{n,m} = \left(-\left(2n + \alpha + \frac{1}{2} \right)^2 + \alpha - \frac{1}{4} \right) f_{n,m}, \tag{75}$$

$$f_{n,m}(t) = {}_4F_3\left(\begin{matrix} -n, & n + \alpha + \frac{1}{2}, & -m - t, & -m + t \\ -m, & \frac{1}{2} - m, & \alpha + 1 & \end{matrix} ; 1 \right),$$

$n = 0, 1, \cdots, m$, we see that in the limit only about half of the spectrum is uncovered. Note that, the functions $f_{n,m}$ are even functions of t whereas $\widehat{f}_{k,m}(-t) = (-1)^k \widehat{f}_{k,m}(t)$.

The recurrences for mutiplication by $2it$ and $-4t^2$ are compatible, so we obtain the following identity relating a special case of Wilson polynomials with a special case of the Wilson polynomials:

$$
{}_4F_3\left(\begin{matrix} -n, & n+\alpha+\frac{1}{2}, & -m-t, & -m+t \\ -m, & \frac{1}{2}-m, & \alpha+1 & \end{matrix}\ ;1\right) = \tag{76}
$$

$$
{}_3F_2\left(\begin{matrix} -2n, 2n+2\alpha+1, -t-m \\ \alpha+1, -2m \end{matrix}\ ;1\right), \quad n = 0, 1, \cdots, m.
$$

6.2 Contraction $E1 \rightarrow E6$

Jacobi → Gegenbauer: System E1 contracts to E6 by setting $a_1 = 0$ and $a_3 = 0$ ($\beta_3 = 0$) to obtain the system E6. The contraction of the operators takes the form

$$
X'^2 = \lim_{\epsilon=0} L_1, \qquad L_1'^2 = \lim_{\epsilon=0} L_2, \qquad L_2' = [L_1, X]. \tag{77}
$$

To investigate the effect of this contraction on the models, we begin with the differential operator model of $E1$ with $a_1 = \beta_1 = 0$ (42), set $\beta_3 = -1/2$ and perform the change of basis (77). The operator X^2 is then given by $X^2 = E'(x+1)/2$, suggesting the change of variables $x = 2t^2 - 1$. Now the model takes the form,

$$
X' = \sqrt{E}t, \quad L_1' = (1-t^2)\partial_t^2 - 2t(1+\beta_2)\partial_t - \beta_2 - \frac{1}{2}, \quad L_2' = [L_1', X]. \tag{78}
$$

The eigenfunctions of L_1' are given by the Gegenbauer polynomials,

$$
C_k^{\beta_2+1/2}(t) = \frac{(2\beta_2+1)_k}{k!}\,{}_2F_1\left(\begin{matrix} -k,\ 2\beta_2+1+k \\ \beta_2+1 \end{matrix}\ \middle|\ \frac{t-1}{2}\right). \tag{79}
$$

The eigenfunction of L_2 in the model for E1, contract to eigenfunctions of L_1'

$$
f_n = {}_2F_1\left(\begin{matrix} -n, & \beta_2+n+1/2 \\ \beta_2+1 & \end{matrix}\ ;t^2-1\right). \tag{80}
$$

The expansion coefficients of the action of X^2 on this basis are

$$
K'(n+1,n) = \frac{2E'(2\beta_2+2n+1)(\beta_2+n+1)}{(2\beta_2+4n+1)(2\beta_2+4n+3)},
$$

$$
K'(n-1,n) = \frac{E'2n(2n-1)}{(2\beta_2+4n-1)(2\beta_2+4n+1)}, \tag{81}
$$

$$
K'(n,n) = E' - K'(n+1,n) - K'(n-1,n) \tag{82}
$$

which suggests half-integer values for n in the model. Indeed, the contraction limit of the basis polynomials gives only half the spectrum. The full spectrum is obtained from eigenfunctions of L'_1 obtained directly as

$$g_k(t) = \frac{k!}{(\beta_2)_k} C_k^{\beta_2/2}(t), \qquad k = 0, 1, 2, \ldots. \tag{83}$$

These polynomials are related to f_n (the contracted basis) as $f_n(t) = g_{2k}(t)$, giving the following identity:

$$ {}_2F_1\left(\begin{matrix} -k, & \beta_2 + k + 1/2 \\ \beta_2 + 1 & \end{matrix} ; t^2 - 1 \right) = {}_2F_1\left(\begin{matrix} -2k,\ 2\beta_2 + 1 + 2k \\ \beta_2 + 1 \end{matrix} \middle| \frac{t-1}{2} \right). $$

6.3 Contraction $E8(\omega = 0) \to E14$:

This contraction leads to Bessel funtions, not to orthogonal polynomials.

7 Degenerate/singular system contractions

7.1 Contraction $S3 \to E3$

Special Dual Hahn → Special Kawtchouk: In model (68) we set $\tilde{a} = \epsilon^2 a$,

$$\begin{pmatrix} \tilde{L}_1 \\ \tilde{L}_2 \\ \tilde{H} \\ \tilde{X} \end{pmatrix} = \begin{pmatrix} \epsilon & 0 & 0 & 0 \\ 0 & \epsilon & 0 & 0 \\ 0 & 0 & \epsilon & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ H \\ X \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ -\epsilon \\ 0 \end{pmatrix} a$$

and obtain in the limit $L'_i = \lim_{\epsilon \to} \tilde{L}_i$ with $a' = -\omega^2$ the model becomes

$$\begin{array}{ll} H' = -2\omega(M+1), & iX' = (s - M)E^1 - sE^{-1}, \\ L'_1 = \omega(2s + 2M + 3), & L'_2 = -\omega(s-M)E^1 - sE^{-1} \end{array} \tag{84}$$

The functions which form the basis for this representation are

$$f_N = (-1)^N {}_2F_1\left(\begin{matrix} -N, & -s \\ -M & \end{matrix} ; 2 \right),$$

special Krawtchouk or Meixner polynomials, depending on whether M is a positive integer. The eigenvalues of iX' are $M, M-2, \cdots, -M$ in the finite case. The action of L'_1 is $L'_1 f_N = (N - M) f_{N+1} - (M+1) f_N - N f_{N-1}$, and the action of L'_2 follows from commutation relation $[L'_1, X'] = 2L'_3$.

For the second model (72), after contraction we have

$$\begin{array}{l} L_1 = \omega\left((x-2m)E + xE^{-1}(-2m-1)\right, \quad X = 2i(x-m), \\ L_2 = -i\omega\left((x-2m)E - xE^{-1}\right) \end{array} . \qquad (85)$$

The eigenvalues of L_1 are $-\omega(2k+1)$, $k = 0, 1, \cdots, 2m$ for finite dimensional representations, and the corresponding eigenfunctions are special:

$$f_k(x) = {}_2F_1\left(\begin{array}{cc} -k, & -x \\ -2m & \end{array} ; 2\right).$$

7.2 Contraction $E1(a_1 = 0) \to$ singular Laguerre:

We use model (42) and let $\beta_2 = 1/\epsilon$, $E' = E/\epsilon$, $u = \epsilon v$, $\epsilon \to 0$. The new basis functions are

$$g_n = {}_1F_1\left(\begin{array}{c} -n \\ \beta_3 + 1 \end{array} ; v\right),$$

associated Laguerre polynomials. The contraction of the operators which corresponds to this limit is

$$L_1' = L_1, \qquad L_2' = \epsilon L_2, \qquad H' = H. \qquad (86)$$

The model contracts to $H' = E'$ and

$$\begin{array}{l} L_1' = 2v, \\ L_2' = \left[4v\partial_{vv} + 4(-v + \beta_3 + 1)\partial_v - 2(\beta_3 + 1)\right]. \end{array} \qquad (87)$$

Whose action on the basis (7.2)

$$\begin{array}{l} L_1' g_n = 2v g_n = -2(\beta_3 + n + 1)g_{n+1} - 2(\beta_3 + 2n + 1)g_n - 2n g_{n-1},, \\ L_2' g_n = \left[4v\partial_{vv} + 4(-v + B_2 + 1)\partial_v - 2(B_2 + 1)\right] g_n. \end{array} \qquad (88)$$

The corresponding limit in the physical system is obtained by taking $y = y'/\sqrt{\epsilon}$, $\beta_2 = 1/\epsilon$ so that the limit corresponds to a subclass of singular quantum systems. Indeed, letting $\epsilon \to 0$: $K = \frac{1}{y'^2}$ a constant that we can set to 1. $S_1 = \partial_{xx} + \frac{\frac{1}{4} - B_2^2}{x^2}$, $S_3 = \partial_{xx} + \frac{\frac{1}{4} - B_2^2}{x^2} - x^2$. Then S_1, S_3 generate the Lie structure algebra $s\ell(2)$.

3) Contraction $E6 \to$ oscillator algebra**:** We use the Gegenbauer model (78) for $E6$ and perform a gauge transformation to get basis functions

$$h_k(v) = \frac{k!}{\sqrt{(2B_1 + 1)_k}} C_k^{B_1 + \frac{1}{2}}(v), \qquad (89)$$

and set $B_1 = 1/\epsilon$, $v = \sqrt{\epsilon}u$, $\sqrt{M} = -i\sqrt{E}/2\sqrt{\epsilon}$. Taking $X = \hat{X} = \sqrt{E}u$, $L_1 = \epsilon\hat{L}_2 = \sqrt{E}(\partial_u - u)$, $L_2 = \epsilon L''_3 = \partial_{uu} - 2u\partial_u - 1$, we find

$$[L_2, X] = 2L_1,\ [L_2, L_1] = 2X,\ [X, L_1] = -E,$$

and Casimir $X^2 - L_1^2 + EL_2 = 0$. In the limit the basis polynomials $\ell_k(u)$ satisfy the recurrence $u\ell_k = \frac{1}{\sqrt{2}}\ell_{k+1} + \frac{k}{\sqrt{2}}\ell_{k-1},\ \ k = 0, 1, \cdots$, which defines the Hermite polynomials $H_k(u)$: $\frac{H_k(u)}{2^{k/2}k!} = \lim_{\lambda\to\infty}\frac{C^{\lambda}(\frac{u}{\sqrt{\lambda}})}{\lambda^{k/2}}$. In the original quantum system we set $x = t/\sqrt{\epsilon}$. Then

$$L_1 = y/t^2, \quad L_2 = t^2\partial_{yy} - y^2/t^2, \quad X = \partial_y, \quad E = -1/t^2.$$

Note that operators X, L_1, L_2, E determine the oscillator algebra, the Lie algebra generated by the annihilation/creation operators for bosons, a, a^*, the number of particles operator $N = a^*a$ and the identity I.

8 Contractions to Laplace Beltrami equations

1) Contraction $E6 \to$ **plane:** We set $a = \frac{1}{4} - B_1^2$, $J = x\partial_y - y\partial_x$ in the $E6$ Hamiltonian. The model is (78) with basis $\{g_k, k = 0, 1, 2. \cdots\}$, and action

$$\hat{X}_2 = 2iMv,\ \hat{L}_2 = (1 - v^2)\partial_{vv} - 2(B_1 + 1)v\partial_v - \frac{1}{2}(2B_1 + 1),$$

$$\hat{L}_1 = -2iM\left((v^2 - 1)\partial_v + (B_1 + 1)v\right), \quad \hat{H} = -4M^2,$$

$$g_k(v) = \frac{k!}{(2B_1 + 1)_k}C_k^{B_1 + \frac{1}{2}}(v), \quad k = 0, 1, \cdots .$$

We contract to the free space Hamiltonian by letting $B_1 \to -1/2$: $\hat{X}_2 g_k = iMg_{k+1} + iMg_{k-1}$, $\hat{L}_2 g_k = -k^2 g_k$. In the limit we find the continuous Tchebicheff polynomials $g_k(x) = 2T_k(x) = k\lim_{s\to 0}\frac{C_k^s(v)}{s}$. After the contraction we have $\hat{L}_2 = \hat{J}^2$, $\hat{J} = \sqrt{1 - v^2}\partial_v$, and $\hat{J}$ generates the new symmetry $\hat{X}_1 = [\hat{J}, \hat{X}_2] = 2iM\sqrt{1 - v^2}$. Since $[\hat{X}_1, \hat{X}_2] = 0$ and $\hat{L}_1 = \frac{1}{2}\{\hat{J}, \hat{X}_1\}$, the quadratic algebra now closes to $e(2)$ the Euclidean Lie algebra.

2) Contraction $S3 \to 2 -$ sphere : We use model (66) . In the finite case the multiplicity of E is $M + 1$. To go to the Laplace-Beltrami eigenvalue equation on the sphere we let $B \to -1/2$. Then cancellation occurs and we have $X = \frac{(s-M)}{2}E^1 - \frac{(s+M)}{2}E^{-1}$, $L_2 = S^2$, $S = is$. From $[X, S]$ we obtain $Y = \frac{(s-M)}{2}E^1 + \frac{(s+M)}{2}E^{-1}$ and these 3 generators define the Lie algebra $so(3)$. With this new symmetry the dimension of the finite representations becomes

$2M + 1$, the basis functions are polynomials in s, rather than s^2 and the spectrum of s is $-M, -M+1, \cdots, M$. The new basis polynomials are

$$ {}_2F_1\left(\begin{array}{cc} -N, & -s-M \\ -2M & \end{array} ; 2 \right), $$

special Krawtchouk polynomials $K_N(s+M; \frac{1}{2}, 2M)$ in the finite dimensional case and special Meixner polynomials $M_N(s+M; -2M, -1)$ in the infinite case.

9 The contraction scheme and final comments

The top half of Figure 1 shows the standard Askey Scheme indicating which orthogonal polynomials can be obtained by pointwise limits from other polynomials and, ultimately, from the Wilson or Racah polynomials. The bottom half of Figure 1 shows how each of the superintegrable systems can be obtained by a series of contractions from the generic system $S9$. Not all possible contractions are listed, partly due to complexity and partly to keep the graph from being too cluttered. (For example, *all* nondegenerate and degenerate superintegrable systems contract to the Euclidean system $H = \partial_{xx} + \partial_{yy}$.) The *singular systems* are superintegrable in the sense that they have 3 algebraically independent generators, but the coefficient matrix of the 2nd order terms in the Hamiltonian is singular. Figure 2 shows which orthogonal polynomials are associated with models of which quantum superintegrable system and how contractions enable us to reach all of these functions from $S9$. Again not all contractions have been exhibited, but enough to demonstrate that the Askey Scheme is a consequence of the contraction structure linking 2nd order quantum superintegrable systems in 2D. It is worth remarking that forthcoming papers by us will simplify considerably the compexity of our approach. We will show that the structure equations for nondegenerate superintegrable systems can be derived directly from the expression for R^2 alone, and the structure equations for degenerate superintegrable systems can be derived, up to a multiplicative factor, from the Casimir alone. It will also be demonstrated that all of the contractions of quadratic algebras in the Askey scheme can be induced by natural contractions of the Lie algebras $e(2, \mathbb{C})$ and $o(3, \mathbb{C})$.

This method obviously extends to 2nd order systems in more variables. A start on this study can be found in [24] (2011). To extend the method to Askey-Wilson polynomials we would need to find appropriate q-quantum mechanical systems with q-symmetry algebras and we have not yet been able to do so.



Askey Scheme of Hypergeometric Orthogonal Polynomials



Partial list of contractions of superintegrable systems

Figure 1: The Askey scheme and contractions of superintegrable systems



Figure 2: The Askey contraction scheme

Acknowledgement

This work was partially supported by a grant from the Simons Foundation (# 208754 to Willard Miller, Jr.).

References

[1] Superintegrability in Classical and Quantum Systems, Tempesta P., Winternitz P., Miller W., Pogosyan G., editors, AMS, vol. 37, 2005.

[2] Kalnins, E. G., Kress J. M. and Miller, W. Jr. Superintegrability and higher order integrals for quantum systems, arXiv:1002.2665v1 [math-ph], J. Phys. A: Math. Theor. 43 (2010) 265205; Structure relations for the symmetry algebras of quantum superintegrable systems, J. Phys.: Conf. Ser. 343 012075 doi:10.1088/1742-6596/343/1/012075 (2012)

[3] Friš J., Mandrosov V., Smorodinsky Ya.A, Uhlír M. and Winternitz P., On Higher Symmetries in Quantum Mechanics; *Phys. Lett.* 1965, V.16, 354–356.

[4] Tremblay F,, Turbiner A. and Winternitz P., An infinite family of solvable and integrable quantum systems in the plane. J.Phys.A: Math. Theor., **42**, (2009) 242001; Periodic orbits for an infinite family of classical superintegrable systems. J. Phys. A: Math. Theor. **43** (2010) 015202.

[5] Wojciechowski S., Superintegrability of the Calogero-Moser System. *Phys. Lett.*, 1983, V. A 95, 279–281.

[6] Evans N.W., Super-Integrability of the Winternitz System; *Phys. Lett.* 1990, V.A 147, 483–486.

[7] Quesne C., Exceptional orthogonal polynomials, exactly solvable potentials and supersymmetry. J. Phys. A: Math. Theor. **41** (2008) 392001; Generalized deformed parafermions, nonlinear deformations of so(3) and exactly solvable potentials, *Phys. Lett. A*, 1994, **193**, 245-250.

[8] Rañada M. F., Superintegrable n=2 systems, quadratic constants of motion, and potentials of Drach. *J. Math. Phys.*, 1997, V.38, 4165–78.

[9] Ballesteras A. and Herranz, F. J., Universal integrals for superintegrable systems on N-dimensional spaces of constant curvature. *J. Phys. A: Math. Theor.* , **40**, F51-F59, (2007).

[10] Fordy, A. P., Quantum Super-Integrable Systems as Exactly Solvable Models , SIGMA **3** 025, 2007.

[11] Kalnins E. G., Kress J .M, and Miller W. Jr., Second order superintegrable systems in conformally flat spaces. I: 2D classical structure theory. *J. Math. Phys.*, 2005, **46**, 053509; Second order superintegrable systems in conformally flat spaces. II: The classical 2D Stäckel transform. *J. Math. Phys.*, 2005, **46**, 053510; Second order superintegrable systems in conformally flat spaces. V: 2D and 3D quantum systems. *J. Math. Phys.*, 2006, **47**, 09350; Nondegenerate 2D complex Euclidean superintegrable systems and algebraic varieties, *J. Phys. A: Math. Theor.*, 2007a, **40**, 3399-3411.

[12] Kress J. M., Equivalence of superintegrable systems in two dimensions. *Phys. Atomic Nuclei*, **70**, 560-566 (2007)

[13] Kalnins E. G., Kress J. M, Miller W. Jr. and Post S., Structure theory for second order 2D superintegrable systems with 1-parameter potentials, *SIGMA*, **5**(2009), 008,

[14] Bonatos D., Daskaloyannis C. and Kokkotas K., Deformed Oscillator Algebras for Two-Dimensional Quantum Superintegrable Systems; *Phys. Rev.*, 1994, V.A 50, 3700–3709.

[15] Letourneau P. and Vinet L., Superintegrable systems: Polynomial Algebras and Quasi-Exactly Solvable Hamiltonians. *Ann. Phys.*, 1995, V.243, 144–168.

[16] Zhedanov A. S., “Hidden symmetry” of Askey-Wilson polynomials, *Theoret. and Math. Phys.*, 1991, **89** , 1146–1157.

[17] Granovskii Ya. I., Zhedanov A. S., and Lutsenko I. M., Quadratic algebras and dynamics in curved spaces. I. Oscillator, *Theoret. and Math. Phys.*, 1992, **91**, 474-480; Quadratic algebras and dynamics in curved spaces. II. The Kepler problem, *Theoret. and Math. Phys.*, 1992, **91**, 604-612.

[18] Daskaloyannis C., Quadratic Poisson algebras of two-dimensional classical superintegrable systems and quadratic associative algebras of quantum superintegrable systems, *J. Math. Phys*, 2001, **42**, 1100-1119, math-ph/0003017.

[19] Daskaloyannis C. and Ypsilantis K., Unified treatment and classification of superintegrable systems with integrals quadratic in momenta on a two dimensional manifold. *J. Math Phys.*, **47**, 042904 (2006) -

[20] Daskaloyannis C. and Tanoudis Y., Quantum superintegrable systems with quadratic integrals on a two dimensional manifold. *J. Math Phys.*, **48**, 072108 (2007)

[21] Koornwinder, T. H., (1988), Group theoretic interpretations of Askey's scheme of hypergeometric orthogonal polynomials, Orthogonal polynomials and their applications (Segovia, 1986), Lecture Notes in Math., 1329, Berlin, New York: Springer-Verlag, pp. 4672, 1988.

[22] Koekoek, R., Lesky, P. A. and Swarttouw, R. F., (2010), Hypergeometric orthogonal polynomials and their q-analogues, Springer Monographs in Mathematics, Berlin, New York: Springer-Verlag, 2010.

[23] Mostafazadeh A., Pseudo-Hermiticity versus PT Symmetry: The necessary condition for the reality of the spectrum of a non-Hermitian Hamiltonian. *J. Math. Phys.* **43**, 205 (2002).

[24] Kalnins E. G., Miller W. Jr and Post S., Wilson polynomials and the generic superintegrable system on the 2-sphere, *J. Phys. A: Math. Theor.* **40**, 11525-11538, (2007), Coupling constant metamorphosis and Nth order symmetries in classical and quantum mechanics, *J. Phys. A: Math. Theor.* **43** (2010) 035202. Two-variable Wilson polynomials and the generic superintegrable system on the 3-sphere, *SIGMA*, **7** (2011), 051, 26 pages

[25] Kalnins E. G., Kress J. M., Miller W. Jr. and Pogosyan G. S., *Completeness of superintegrability in two-dimensional constant curvature spaces. J. Phys. A: Math Gen.*, 2001, V.34, 4705–4720.

[26] Kalnins E. G., Miller W. Jr., Subag E and Heinenen R., Contractions of 2nd order superintegrable systems in 2D. (In preparation). (2012).

[27] Inönü, E. and Wigner, E. P., On the contraction of groups and their representations. *Proc. Nat. Acad. Sci. (US) (***39**, 510-524, (1953).

[28] Andrews G. E., Askey, R. and Roy R., Special Functions. Encyclopedia of mathematics and its Applications, *Cambridge University Press*, Cambridge, UK, 1999.

[29] Askey, R., An integral of Ramanujan and orthogonal polynomials. T*he Journal of the Indian Mathematical Society.* New Series **51**, 2736, (1987).

[30] Kalnins E. G., Miller W. Jr., Williams G. C. and Pogosyan G. S., On superintegrable symmetry-breaking potentials in n-dimensional Euclidean space. *J. Phys. A: Math Gen.*, 2002, V.35, 4655–4720.

[31] Kalnins E. G., Kress J. M., Miller, W. Jr. and Winternitz P., *Superintegrable systems in Darboux spaces. J. Math. Phys.*, 2003, V.44, 5811–5848.

[32] Krall H. L. and Frink O., A new class of orthogonal polynomials: The Bessel polynomials. *Trans. Amer. Math. Soc.*, **63**, (1949) pp. 100115